

# TransIT<sup>®</sup>-3T3 Transfection Kit

Protocol for MIR 2180, 2184, 2185, 2186



The Transfection Experts

## INTRODUCTION

TransIT<sup>®</sup>-3T3 Transfection Kit is specifically optimized to provide exceptional transfection efficiency of plasmid DNA in NIH 3T3 cells and cell types of associated lineage. TransIT-3T3 Transfection Kit contains two components, namely: TransIT<sup>®</sup>-3T3 Reagent and the 3T3 Authority Reagent. This kit provides all the attributes of the trusted TransIT series of transfection reagents: high transfection efficiency, low toxicity, serum compatibility, simplicity of use and reproducibility. Transfection using TransIT-3T3 Transfection Kit does not require medium changes and can be carried out in serum-containing medium. This kit is suitable for both transient and stable transfection.

## SPECIFICATIONS

<b>Storage</b>	Store both TransIT-3T3 Reagent and 3T3 Authority Reagent at -20°C. <b>Before each use</b> , warm to room temperature and vortex gently.
<b>Product Guarantee</b>	6 months from the date of purchase, when properly stored and handled.



Warm TransIT-3T3 and 3T3 Authority Reagent to room temperature and vortex gently before each use.

## MATERIALS

### Materials Supplied

TransIT-3T3 Transfection Kit is supplied in *one* of the following formats.

Product No.	Volume of TransIT-3T3 Reagent	Volume of 3T3 Authority Reagent
MIR 2184	1 × 0.4 ml	1 × 0.2 ml
MIR 2180	1 × 1 ml	1 × 0.5 ml
MIR 2185	5 × 1 ml	5 × 0.5 ml
MIR 2186	10 × 1 ml	10 × 0.5 ml

### Materials required, but not supplied

- Cultured 3T3 cells
- Appropriate cell culture medium
- Purified DNA
- Serum-free medium (e.g. Opti-MEM<sup>®</sup> I Reduced-Serum Medium)
- Sterile tube for transfection complex preparation
- Micropipets
- Reporter assay as required

**For Research Use Only.**

## BEFORE YOU START:

### Important Tips for Optimal DNA Transfection

Optimize reaction conditions for each 3T3 cell subtype to ensure successful transfections. The suggestions below yield high efficiency DNA transfection using the *TransIT*-3T3 Transfection Kit.

**Table 1** on Page 3 presents recommended starting conditions depending on culture vessel size.

- **Cell density (% confluence) at transfection.** The recommended cell density for 3T3 cell subtypes at transfection is 50–80% confluence. Determine the optimal cell density for each 3T3 cell subtype in order to maximize transfection efficiency. Divide the cells 18–24 hours before transfection to ensure that the cells are actively dividing and reach the appropriate cell density at the time of transfection.
- **DNA purity.** Use highly purified, sterile, and contaminant-free DNA for transfection. Plasmid DNA preps that are endotoxin-free and have  $A_{260/280}$  absorbance ratio of 1.8–2.0 are desirable. DNA prepared using miniprep kits is not recommended as it may contain high levels of endotoxin. We recommend using MiraCLEAN<sup>®</sup> Endotoxin Removal Kit (MIR 5900) to remove any traces of endotoxin from your DNA preparation.
- ***TransIT*-3T3 Reagent:DNA ratio.** As a starting point, use 3  $\mu$ l of *TransIT*-3T3 Reagent per 1  $\mu$ g of DNA. The optimal *TransIT*-3T3 Reagent to DNA ratio can be determined by titrating the reagent from 2–6  $\mu$ l per  $\mu$ g of DNA. Please refer to **Table 1** on Page 3 for recommended starting conditions.
- **3T3 Authority Reagent:DNA ratio.** As a starting point, use 1  $\mu$ l of 3T3 Authority Reagent per 1  $\mu$ g of DNA. The optimal 3T3 Authority Reagent:DNA ratio should be determined by titrating the reagent from 0.5–3  $\mu$ l per  $\mu$ g of DNA. Refer to **Table 1** on Page 3 for recommended starting conditions.
- **Complex formation conditions.** Prepare *TransIT*-3T3:3T3 Authority:DNA complexes in serum-free growth medium. Mirus recommends Opti-MEM I Reduced-Serum Medium.
- **Cell culture conditions:** Culture cells in the appropriate medium, with or without serum. There is no need to perform a medium change to remove the transfection complexes. *TransIT*-3T3 Transfection Kit yields improved efficiencies when transfections are performed in complete growth medium (instead of serum-free medium) without a post-transfection medium change.
- **Presence of antibiotics:** Antibiotics will inhibit transfection complex formation and therefore should be excluded from the complex formation step. Transfection complexes can be added to cells grown in complete culture medium containing low levels of antibiotics (0.1–1X final concentration of penicillin/streptomycin mixture).
- **Post-transfection incubation time.** Determine the best incubation time post-transfection for each cell type. The optimal incubation time is generally 24–72 hours, but will vary depending on the goal of the experiment, nature of the plasmid, and the half-life of the expressed protein.



**Do not** use DNA prepared using miniprep kits for transfection.



**Do not** use serum or antibiotics in the medium during transfection complex formation.

## DNA TRANSFECTION PROTOCOL

The following procedure describes how to perform DNA transfections in 6-well plates. The surface areas of other culture vessels are different and transfections must be scaled accordingly. Appropriately increase or decrease the amounts of serum free medium, *TransIT*-3T3 Reagent, 3T3 Authority Reagent, DNA and complete culture medium based on the surface area of the cell culture vessel (please refer to **Table 1** on Page 3).

**Table 1.** Recommended starting conditions for DNA transfections with the *TransIT*<sup>®</sup>-3T3 Transfection Kit.

Culture vessel	96-well plate	48-well plate	24-well plate	12-well plate	6-well plate	10-cm dish	T75 flask
Surface area	0.35 cm <sup>2</sup>	1.0 cm <sup>2</sup>	1.9 cm <sup>2</sup>	3.8 cm <sup>2</sup>	9.6 cm <sup>2</sup>	59 cm <sup>2</sup>	75 cm <sup>2</sup>
Complete growth medium	92 µl	263 µl	0.5 ml	1.0 ml	2.5 ml	15.5 ml	19.7 ml
Serum-free medium	9 µl	26 µl	50 µl	100 µl	250 µl	1.5 ml	1.9 ml
DNA (1µg/µl stock)	0.09 µl	0.25 µl	0.5 µl	1 µl	2.5 µl	15.5 µl	19.7 µl
<i>TransIT</i> -3T3 Reagent	0.27 µl	0.75 µl	1.5 µl	3 µl	7.5 µl	46.5 µl	59.1 µl
3T3 Authority Reagent	0.09 µl	0.25 µl	0.5 µl	1 µl	2.5 µl	15.5 µl	19.7 µl



Surface areas are based on Greiner tissue culture plates and Falcon 10-cm dishes and T75 flasks. All volumes given are per well (or per dish) for a given culture vessel.

If small volumes of reagents need to be pipetted, dilute the *TransIT*-3T3 reagent in 80% ethanol. The 3T3 Authority reagent can be diluted in sterile water. **Do not** store diluted reagents.

### Transient DNA transfection protocol per well of a 6-well plate

#### A. Plate cells

1. Approximately 18–24 hours before transfection, plate cells in 2.5 ml complete growth medium per well in a 6-well plate. Ideally cells should be 50–80% confluent (~1–3 × 10<sup>5</sup> cells/well) prior to transfection.
2. Incubate the cell cultures overnight.

#### B. Prepare *TransIT*-3T3:3T3 Authority:DNA complexes (Immediately before transfection)

1. Warm *TransIT*-3T3 and 3T3 Authority reagents to room temperature and vortex gently before using.
2. Place 250 µl of Opti-MEM I Reduced-Serum Medium in a sterile tube.
3. Add 2.5 µg (2.5 µl of a 1 µg/µl stock) DNA. Pipet gently to mix completely.
4. Add 7.5 µl *TransIT*-3T3 Reagent to the diluted DNA mixture. Pipet gently to mix completely.
5. Add 2.5 µl 3T3 Authority Reagent to the diluted *TransIT*-3T3:DNA mixture. Pipet gently to mix completely.
6. Incubate at room temperature for 15–30 minutes.



Divide cultured cells 18–24 hours before transfection to ensure active cell division at the time of transfection.



Warm *TransIT*-3T3 and 3T3 Authority reagents to room temperature and vortex gently before each use.

**C. Distribute the complexes to cells in complete growth medium**

1. Add the complexes drop-wise to different areas of the wells. It is not necessary to replace the complete growth medium with fresh medium.
2. Gently rock the culture vessel back-and-forth and from side-to-side to evenly distribute the *TransIT*-3T3:3T3 Authority:DNA complexes.
3. Incubate for 24–72 hours. It is not necessary to replace the complete growth medium with fresh medium.
4. Harvest cells and assay as required.



There is no need to change fresh culture medium after transfection. If required, perform a medium change at least 4 hours post-transfection.

For generating stable cell transfectants, passage the cells 24–48 hours post-transfection in complete growth medium containing the appropriate selection antibiotic such as G418 or Hygromycin B. Maintain selection for 1–2 weeks, allowing selection of cells that have undergone stable integration of DNA.

## TROUBLESHOOTING GUIDE

Problem	Solution
<b>LOW DNA TRANSFECTION EFFICIENCY</b>	
<i>TransIT</i> -3T3 or 3T3 Authority Reagent was not mixed properly.	Warm <i>TransIT</i> -3T3 and 3T3 Authority Reagents to room temperature and vortex gently before each use.
Suboptimal amount of <i>TransIT</i> -3T3 Reagent	Determine optimal amount of <i>TransIT</i> -3T3 Reagent for each 3T3 cell subtype. Titrate the <i>TransIT</i> -3T3 Reagent from 2–6 µl per 1 µg DNA. Refer to “Before You Start” on Page 2.
Suboptimal amount of 3T3 Authority Reagent	Determine optimal amount of 3T3 Authority Reagent for each 3T3 cell subtype. Titrate the 3T3 Authority Reagent from 0.5–3 µl per 1 µg DNA. Refer to “Before You Start” on Page 2.
Suboptimal DNA concentration	<p>Confirm DNA concentration and purity. Use plasmid DNA preps that have an <math>A_{260/280}</math> absorbance ratio of 1.8–2.0.</p> <p>The optimal DNA concentration generally ranges between 1–3 µg/well of a 6-well plate. Start with 2.5 µg/well of a 6-well plate. Consider testing more or less DNA while scaling the amount of <i>TransIT</i>-3T3 and 3T3 Authority Reagents accordingly.</p>
Low-quality plasmid DNA	<p>Use highly purified, sterile, endotoxin and contaminant-free DNA for transfection.</p> <p>We recommend using Mirus Bio’s MiraCLEAN® Endotoxin Removal Kit (MIR 5900) for removal of endotoxin from your DNA preparation.</p> <p>Alternatively, use cesium chloride gradient or anion exchange purified DNA which contains levels of endotoxin that do not harm most cells.</p> <p><b>Do not</b> use DNA prepared using miniprep kits as it may contain high levels of endotoxin.</p>
Inhibitor present during transfection	<p>Serum and antibiotics inhibit transfection complex formation. Prepare transfection complexes in serum-free growth medium. We recommend Opti-MEMI Reduced-Serum Medium. Once transfection complexes are formed, they can be added directly to cells cultured in complete growth medium containing serum and 0.1–1X antibiotics.</p> <p>Polyanions such as dextran sulfate or heparin can inhibit transfection. Use culture medium that does not contain these polyanions. If necessary, the transfection medium can be replaced with polyanion containing medium 24 hours post transfection.</p>
Incorrect vector sequence	If you do not observe expression of your target insert, verify the sequence of the plasmid DNA.
Transfection incubation time	Determine the optimal transfection incubation time for each cell type and experiment. Test a range of incubation times (e.g. 12–72 hours). The best incubation time is generally 24–48 hours.
Cells not actively dividing at the time of transfection	Divide the culture at least 18–24 hours before transfection to ensure that the cells are actively dividing and reach optimal cell density at time of transfection.
Precipitate formation during transfection complex formation	<p>During complex formation, scale all reagents including serum-free medium, <i>TransIT</i>-3T3 Reagent, 3T3 Authority Reagent and plasmid DNA according to Table 1 on Page 3.</p> <p>Precipitation may be observed when excess DNA is used during complex formation. This may negatively impact transfection efficiency. To avoid precipitation when using high concentrations of DNA, increase the volume of serum-free medium during complex formation by two-fold.</p>

## TROUBLESHOOTING GUIDE continued

Problem	Solution
<b>LOW DNA TRANSFECTION EFFICIENCY</b>	
Proper experimental controls were not included	To verify efficient transfection, use <i>TransIT-3T3</i> Transfection Kit to deliver a positive control such as a luciferase, beta-galactosidase or green fluorescent protein (GFP) encoding plasmid. To assess delivery efficiency of plasmid DNA, use Mirus' <i>Label IT<sup>®</sup> Tracker™</i> Intracellular Nucleic Acid Localization Kit to label the target plasmid or Mirus' pre-labeled <i>Label IT<sup>®</sup> Plasmid Delivery Controls</i> (please refer to Related Products on Page 7).
<b>HIGH CELLULAR TOXICITY</b>	
Transfection complexes and cells not mixed thoroughly after complex addition	Add transfection complexes drop-wise to different areas of the wells containing plated cells . Gently rock the dish back-and-forth and from side-to-side to distribute the complexes evenly. Do not swirl or rotate the dish, as this may cause uneven distribution.
Transfection complexes added to cells cultured in serum-free medium	Allow transfection complexes to form in serum-free medium, then add these complexes to cells cultured in complete growth medium. The presence of serum in the growth medium improves transfection efficiency and reduces cytotoxicity. No culture medium change is required after the addition of transfection complexes to cells.
Endotoxin-contaminated plasmid DNA	Use highly purified, sterile, endotoxin and contaminant-free DNA for transfection.
	We recommend using Mirus Bio's <i>MiraCLEAN<sup>®</sup> Endotoxin Removal Kit (MIR 5900)</i> for removal of any traces of endotoxin from your DNA preparation. Alternatively, use cesium chloride gradient or anion exchange purified DNA which contains levels of endotoxin that do not harm most cells.
Expressed target gene is toxic to cells	Do not use DNA prepared using miniprep kits as it might contain high levels of endotoxin.
	Compare toxicity levels against a cells alone control and cells transfected with an empty vector to assess the cytotoxic effects of the target protein being expressed. If lower levels of target gene expression are desired in your transfection experiments, consider reducing the amount of target plasmid. Maintain the optimal <i>TransIT-3T3:3T3 Authority:DNA</i> ratio by using carrier DNA such as an empty cloning vector.
Cell density not optimal at time of transfection	Determine the best cell density for each 3T3 cell subtype to maximize transfection efficiency. Use this cell density in subsequent experiments to ensure reproducibility. For most 3T3 cell subtypes, 50–80% confluence is recommended at transfection, but use of higher or lower densities may increase cell viability depending on cell type.
Cell morphology has changed	Mycoplasma contamination can alter cell morphology and affect transfection efficiency. Check your cells for Mycoplasma contamination. Use a fresh frozen stock of cells or use appropriate antibiotics to eliminate Mycoplasma.
	A high or low cell passage number can make cells more sensitive and refractory to transfection. Maintain a similar passage number between experiments to ensure reproducibility.

## RELATED PRODUCTS

- Ingenio<sup>®</sup> Electroporation Solution and Kits
- Label IT<sup>®</sup> Plasmid Delivery Controls
- Label IT<sup>®</sup> Tracker™ Intracellular Nucleic Acid Localization Kits
- MiraCLEAN<sup>®</sup> Endotoxin Removal Kits
- TransIT<sup>®</sup>-3D Transfection Reagent
- TransIT<sup>®</sup>-2020 Transfection Reagent
- TransIT-PRO<sup>®</sup> Transfection Kit
- TransIT<sup>®</sup>-LT1 Transfection Reagent
- TransIT<sup>®</sup> Cell Line Specific Transfection Reagents and Kits
- TransIT<sup>®</sup>-QR and TransIT<sup>®</sup>-EE Delivery Solutions and Kits



### Reagent Agent<sup>®</sup>

Reagent Agent<sup>®</sup> is an online tool designed to help determine the best solution for nucleic acid delivery based on in-house data, customer feedback and citations.

Learn more at:  
[www.TheTransfectionExperts.com/reagentagent](http://www.TheTransfectionExperts.com/reagentagent)

For details on our products, visit [www.mirusbio.com](http://www.mirusbio.com) or [www.TheTransfectionExperts.com](http://www.TheTransfectionExperts.com).

Contact Mirus Bio for additional information.



Mirus Bio LLC  
545 Science Drive  
Madison, WI 53711  
Toll-free: 888.530.0801  
Direct: 608.441.2852  
Fax: 608.441.2849

©1996-2012. All rights reserved. Mirus Bio LLC. All trademarks are the property of their respective owners. HeLaMONSTER, Ingenio, Label IT, MiraCLEAN,  $\mu$ Array, pLIVE, siQUEST, TransIT, TransIT-Neural, TransIT-PRO and TransIT-TKO are registered trademarks of Mirus Bio LLC. Tracker is a trademark of Mirus Bio LLC.

Use of Mirus Bio TransIT<sup>®</sup> polyamine transfection reagents are covered by U.S. Patent No. 5,744,335, No. 6,180,784, No. 7,101,995, No. 7,601,367 and patents pending. The use of certain Mirus Bio transfection products are the subject of one or more of U.S. Patents No. 7,335,509, No. 7,655,468 and/or other pending U.S. patent applications. Mirus Bio Label IT<sup>®</sup> nucleic acid labeling and modifying reagents are covered by U.S. Patent No. 6,262,252, No. 6,593,465, No. 7,049,142, No. 7,326,780 and No. 7,491,538. Cy<sup>TM3</sup> and Cy<sup>TM5</sup> products or portions thereof are manufactured under license from Carnegie Mellon University and are covered by U.S. Patent No. 5,268,486.

**This product is sold to the Buyer with a limited license for Research Use Only and is not for clinical, therapeutic or diagnostic use in humans or animals. This product, or parts from this product, may not be re-packaged or re-sold without written permission from Mirus Bio LLC. The buyer agrees not to infringe upon Mirus patents or to attempt to reverse engineer, reconstruct, synthesize or otherwise modify Mirus Bio products. A license from Mirus Bio LLC is required for commercial application of this product. For obtaining a license to use this product for commercial application, contact Mirus Bio LLC, 545 Science Drive, Madison, WI 53711. Email [license@mirusbio.com](mailto:license@mirusbio.com). For full terms and conditions, visit [www.mirusbio.com](http://www.mirusbio.com).**

For publications citing the use of TransIT<sup>®</sup>-3T3 Transfection Kit, visit [www.mirusbio.com](http://www.mirusbio.com).