

# VirusGEN<sup>®</sup> SELECT AAV Transfection Kit



## Protocol for MIR 6770 and MIR 6775

SDS available at [mirusbio.com/6770](http://mirusbio.com/6770)

## INTRODUCTION

Adeno-associated virus (AAV) is a nonenveloped, single-stranded DNA virus from the *Parvoviridae* family notable for its lack of pathogenicity, low immunogenicity and ability to infect both dividing and quiescent cells. Because AAV is replication-defective in the absence of adeno or helper proteins and is not implicated in any known human diseases, it is widely considered a safe gene delivery vehicle for *in vivo* and *in vitro* applications. Accordingly, recombinant AAV has become an invaluable tool for gene therapy and the creation of isogenic human disease models.

Efficient, high titer and large-scale viral vector manufacturing processes are necessary for production of AAV-based therapies. Additionally, raw or ancillary materials used for viral vector manufacturing must be carefully selected as part of a risk-based approach for the development of therapeutics. The *TransIT-VirusGEN<sup>®</sup> SELECT* Transfection Reagent addresses these needs by providing robust titers for AAV production and expert support for researchers developing biotherapeutics. The *VirusGEN<sup>®</sup> SELECT AAV Transfection Kit* further enhances the performance of *TransIT-VirusGEN<sup>®</sup> SELECT* Transfection Reagent in suspension HEK 293 cells through the addition of the proprietary *VirusGEN<sup>®</sup> SELECT AAV Complex Formation Solution and Enhancer*. *TransIT-VirusGEN<sup>®</sup> SELECT* Transfection Reagent is assayed for formulation identity, appearance, sterility, bacterial endotoxin and mycoplasma. In addition to these tests, *VirusGEN<sup>®</sup> SELECT AAV Complex Formation Solution and Enhancer* is assayed for pH and osmolality. Thus, *TransIT-VirusGEN<sup>®</sup> SELECT AAV Transfection Kit* is ideal for generating large-scale AAV preparations with the dependability and quality that are essential for streamlined viral vector production.

## SPECIFICATIONS

<b>Storage</b>	Store <i>TransIT-VirusGEN<sup>®</sup> SELECT</i> Reagent at -10 to -30°C, tightly capped. Store <i>VirusGEN<sup>®</sup> SELECT AAV Complex Formation Solution and Enhancer</i> at 2 to 10°C. <i>Before each use</i> , warm to room temperature and mix gently.
<b>Stability / Guarantee</b>	Guaranteed as noted on the Certificate of Analysis when properly stored and handled.



Warm *TransIT-VirusGEN<sup>®</sup> SELECT* Reagent and *VirusGEN<sup>®</sup> SELECT AAV Complex Formation Solution and Enhancer* to room temperature before each use. Mix gently.

## MATERIALS

### Materials Supplied

The *VirusGEN<sup>®</sup> SELECT AAV Transfection Kit* is supplied in *one* of the following formats.

Product No.	Volume of <i>TransIT-VirusGEN<sup>®</sup> SELECT</i> Transfection Reagent	Volume of <i>VirusGEN<sup>®</sup> SELECT AAV Complex Formation Solution and Enhancer</i>
MIR 6770	1 × 30 ml	1 × 1 L
MIR 6775	1 × 150 ml	5 × 1 L

### For Materials Required but Not Supplied, See Protocol Sections:

- (I) AAV Generation in Suspension HEK 293 Cell Cultures
- (II) AAV Transduction/Titering Method Using a GFP Reporter Virus

***For Research Use Only***

## BEFORE YOU START:

### Important Tips for Optimal AAV Production

Mirus recommends using suspension Expi293F™ Cells (Gibco Cat. No. A14527) for high titer AAV production. The suggestions below yield high efficiency plasmid DNA transfection using the VirusGEN® SELECT AAV Transfection Kit.

- **Cell density at transfection.** The recommended cell density for suspension Expi293F™ cells is  $2 \times 10^6$  cells/ml. Passage cells 18-24 hours before transfection to ensure that cells are actively dividing and reach the appropriate density at time of transfection.
- **DNA purity.** Use highly purified, sterile, endotoxin-free and contaminant-free DNA for transfection. Plasmid DNA preparations that have an  $A_{260/280}$  absorbance ratio of 1.8-2.0 are desirable. DNA prepared using miniprep kits is not recommended as it might contain high levels of endotoxin. We recommend using MiraCLEAN® Endotoxin Removal Kit (MIR 5900) to remove endotoxin from your DNA preparation.
- **AAV packaging and transfer plasmids.** The VirusGEN® SELECT AAV Transfection Kit was optimized using a 1:1:1 weight ratio of pAAV-hrGFP, pAAV-RC and pHelper (AAV Helper-Free System, Agilent Technologies). The optimal ratio may vary for other packaging systems.
- **Ratio of TransIT-VirusGEN® SELECT to DNA.** Determine the optimal TransIT-VirusGEN® SELECT Reagent:DNA ratio for each cell type by varying the amount of reagent from 1.5-3  $\mu$ l per 1-2  $\mu$ g total DNA. Refer to **Table 1** for recommended starting conditions based on culture size.
- **Complex formation conditions.** Prepare TransIT-VirusGEN® SELECT Reagent:DNA complexes in VirusGEN® SELECT AAV Complex Formation Solution and Enhancer. We recommend forming complexes in a volume that is 10% of the total culture volume. If forming complexes in a smaller or larger volume, complex formation time may need to be shortened or extended. We recommend an initial complex formation time of 30 minutes (15-60 minutes yields similar results).
- **Cell culture conditions.** Culture cells in the appropriate complete growth medium, (e.g. Expi293™ Expression Medium). Ensure cells are  $\geq 95\%$  viable by trypan blue exclusion and doubling every 24 hours. After transfection, there is no need to perform a medium change to remove the transfection complexes.
- **Presence of antibiotics.** Antibiotics inhibit transfection complex formation and should be excluded from the complex formation step. Transfection complexes can be added directly to cells growing in complete culture medium containing serum and low levels of antibiotics (0.1-1X final concentration of penicillin/streptomycin mixture).
- **Media change post-transfection.** A media change is not required and could be detrimental to virus titers; therefore, we do not recommend a media change post-transfection.
- **Post-transfection incubation time.** The optimal incubation time for harvesting high titer AAV generated with this kit is generally 48-72 hours post-transfection.




*Premix packaging and transfer plasmids* together prior to adding to the complex formation medium.



*Do not* use serum or antibiotics in the media during transfection complex formation.

Transfection complexes can be added directly to cells cultured in growth media +/- serum and up to 0.1-1X antibiotics.

### VirusGEN® SELECT AAV Transfection Kit Workflow:



## SECTION I: AAV Generation in Suspension HEK 293 Cell Cultures

The following procedure describes plasmid DNA transfections for AAV generation in 125 ml Erlenmeyer shake flasks using 25 ml of complete growth medium. If using an alternate cell culture vessel, increase or decrease the amounts of VirusGEN® SELECT AAV Complex Formation Solution and Enhancer, *TransIT-VirusGEN® SELECT* Reagent and total DNA based on the **volume of complete growth medium** to be used. To calculate the required reagent quantities based on the recommended starting conditions and total culture volume, refer to the calculation worksheet in **Table 1** (below).

**NOTE:** Use of the VirusGEN® SELECT AAV Complex Formation Solution and Enhancer is only recommended for AAV production in suspension HEK 293 cell lines.

**Table 1.** Calculation worksheet for scaling the VirusGEN® SELECT AAV Transfection Kit

Starting conditions per milliliter of complete growth medium				
	Per 1 ml		Total culture volume	Reagent quantities
VirusGEN® SELECT AAV Complex Formation Solution and Enhancer	0.1 ml	×	_____ ml	= _____ ml
Total Plasmid DNA (1 µg/µl stock)	2 µl	×	_____ ml	= _____ µl
<i>TransIT-VirusGEN® SELECT</i> Reagent	3 µl	×	_____ ml	= _____ µl

**NOTE:** Total Plasmid DNA refers to the combined weight of AAV plasmids (in µg) per transfection.



Complexes can also be formed in PBS or serum-free basal medium instead of the VirusGEN® SELECT AAV Complex Formation Solution and Enhancer. However, doing so may result in decreased AAV titer.

### Materials Required but Not Supplied

- Suspension HEK 293 Cells (e.g. Expi293F™ Cells, Gibco Cat. No. A14527)
- Complete Culture Medium (e.g. Expi293™ Expression Medium, Gibco Cat. No. A1435101)
- Plasmid DNA (e.g. Agilent AAV2 pDNA: pAAV-hrGFP (Cat. No. 240074-51), pAAV-RC (Cat. No. 240071-53), pHelper (Cat. No. 240071-54))
- Phosphate Buffered Saline (PBS) (e.g. Millipore Sigma Cat. No. D8537)
- Erlenmeyer shake flasks (e.g. Corning® Cat. No. 431143 or Thomson Cat. No. 931110)
- 10X Cell Lysis Buffer (500 mM Tris pH 8, 10% Tween® 20, 20 mM MgCl<sub>2</sub>)
- 5 M Sodium Chloride (5 M NaCl)
- Benzonase® (e.g. Sigma Cat. No. E1014 or Syd Labs Cat. No. BP4200)
- Reporter assay as required

### Transient Plasmid Transfection Protocol per 25 ml HEK 293 Culture

#### A. Maintenance of cells

1. Passage suspension HEK 293 cells 18-24 hours prior to transfection to ensure that cells are actively dividing at the time of transfection and to obtain a density of  $2 - 4 \times 10^6$  cells/ml the next day.

NOTE: Perform cell counts and evaluate viability daily to ensure that cells are doubling every 24 hours and  $\geq 95\%$  viable by trypan blue exclusion. DO NOT proceed with transfection if cells are not doubling normally or are  $< 95\%$  viable.

2. Incubate cells overnight at appropriate temperature and CO<sub>2</sub> levels (e.g. 37°C, 5-8% CO<sub>2</sub>, shaking).

#### B. Prepare *TransIT-VirusGEN® SELECT*:DNA complexes (immediately before transfection)

1. Immediately prior to transfection, seed cells at a density of  $2 \times 10^6$  cells/ml into a transfection culture vessel (e.g. 25 ml per 125 ml Erlenmeyer shake flask).
2. Warm *TransIT-VirusGEN® SELECT* to room temperature and vortex gently before using.
3. Place 2.5 ml of *VirusGEN® SELECT* AAV Complex Formation Solution and Enhancer in a sterile tube.

NOTE: When evaluating new AAV serotypes or plasmid constructs, Mirus recommends comparing +/- Enhancer, i.e. forming complexes in 2.5 ml of PBS.

4. In a separate sterile tube, combine AAV plasmids per manufacturer recommendations to a final concentration of 1 µg/µl. Mix thoroughly.
5. Transfer 50 µl of the DNA mixture prepared in Step B.4 to the tube containing *VirusGEN® SELECT* AAV Complex Formation Solution and Enhancer. Mix completely.
6. Add 75 µl *TransIT-VirusGEN® SELECT* Reagent to the diluted DNA. Mix completely by inversion or vortexing. Do NOT agitate Reagent:DNA complexes again after this initial mixing.

NOTE: This is a 1.5:1 mixture of transfection reagent to total DNA (vol:wt), which can be further optimized for AAV production using *TransIT-VirusGEN® SELECT* Reagent.

7. Incubate at room temperature for 15-60 minutes without additional agitation to allow transfection complexes to form.

#### C. Distribute the complexes to cells in complete growth medium

1. Add the *TransIT-VirusGEN® SELECT*:DNA complexes (prepared in Step B) to culture vessel, swirling gently to distribute.
2. Shake flasks on an orbital shaker (125 rpm when using a shaker with a 1.9 cm orbital throw) at appropriate temperature and CO<sub>2</sub> levels (e.g. 37°C, 5-8% CO<sub>2</sub>).
3. Incubate cultures for 48-72 hours prior to AAV harvest.

#### D. Harvest and storage of AAV

1. Following the 48-72 hour incubation, transfer the total volume of cell suspension (i.e. 27.5 ml) to a sterile conical tube or appropriate vessel.

NOTE: See 'Freeze/thaw Procedure' below for an alternative method to harvest AAV.

2. Add 0.1X volume of 10X Cell Lysis Buffer (i.e. 2.75 ml) and 100 U/ml Benzonase® (i.e. 2,750 U). Mix completely and incubate at 37°C for 1.5 hours with shaking.
3. Add 0.1X volume of 5 M NaCl (i.e. 2.75 ml) and mix completely. Incubate at 37°C for 30 minutes with shaking.
4. Centrifuge the mixture at  $4,100 \times g$  for 10 minutes to remove cell debris. Carefully transfer the AAV containing supernatant to a new sterile tube.
5. Store AAV stocks at -80°C.



Divide cultured cells 18-24 hours before transfection to ensure that cells are actively dividing at the time of transfection.



Do NOT allow the *TransIT-VirusGEN® SELECT* Reagent to incubate alone in complex formation solution  $> 5$  min, i.e. if the reagent is pre-diluted, add DNA within 5 min for optimal complex formation.

Do NOT agitate Reagent:DNA complexes after the initial mixing. This will result in decreased titer.



There is no need to change culture medium after transfection, unless required by your cell type or culture conditions.



Benzonase® is a non-specific endonuclease used to liberate virus particles from residual nucleic acids in the cell lysates and increase AAV titers.

**Harvest of AAV with Freeze/thaw Procedure (Alternative)**

1. Prepare a dry ice/ethanol bath.
2. Centrifuge cell suspension at  $1,750 \times g$  for 10 minutes.
3. Remove the supernatant and process as needed.
4. Add 5.5 ml of Freeze/Thaw Lysis Buffer (50 mM Tris pH 8.5, 150 mM NaCl, 2 mM  $MgCl_2$ ) to the cell pellet. Mix thoroughly until cell clumps are no longer visible.  
NOTE: The required volume of Freeze/Thaw Lysis Buffer is calculated by multiplying the transfected cell culture volume by 0.2 ml.
5. To ensure sufficient lysis of the cells, freeze cells completely in the dry ice/ethanol bath. Visually inspect to verify that cells are frozen solid, then thaw in a  $37^\circ C$  water bath. Repeat freeze/thaw procedure a total of three times.
6. Add 50 U/ml Benzonase® (i.e. 275 U) to the cell lysate and mix gently but thoroughly. Incubate at room temperature for 30 minutes.
7. Centrifuge the cell lysate at  $10,000 \times g$  for 10 minutes to remove cell debris. Carefully transfer the AAV containing supernatant to a new sterile tube.
8. Store AAV stocks at  $-80^\circ C$ .

## SECTION II: AAV Transduction/Titering Method Using a GFP Reporter Virus

The following procedure describes transduction of HT-1080 cells grown in a 24-well format with a GFP reporter AAV2 and is meant to determine functional AAV2 titers. The number of wells needed for this assay will depend on the number of AAV2 stocks titered and the number of dilutions required for testing per stock (see Step B.2). Testing several dilutions is recommended to accurately determine the functional AAV2 titer. This protocol can be adapted to transduce cells that are permissive to different AAV serotypes.

### Materials Required, but Not Supplied

- HT-1080 cells (ATCC Cat. No. CCL-121)
- Dulbecco's Modified Eagle Medium (DMEM) (Corning Cat. No. 10-013-CV)
- Complete HT-1080 cell culture medium (e.g. DMEM + 10% FBS)
- DMEM + 2% FBS for AAV dilutions
- AAV stock(s) expressing GFP reporter
- 24-well tissue culture plate(s)
- 1X PBS and trypsin
- Flow cytometer equipped with a GFP compatible laser

#### A. Plate cells

1. Approximately 4-6 hours before transduction, plate HT-1080 cells in 0.5 ml complete growth medium per well in a 24-well plate. A starting cell density of  $1 \times 10^5$  cells/ml is recommended. Cells should be adhered to the plate and 40-50% confluent at the time of transduction.
2. Record the cell count, which is critical to determine an accurate functional titer.

#### B. Transduce with GFP-encoding recombinant AAV

1. Thaw AAV stock(s) in 37°C water bath. Remove promptly after virus has thawed to prevent virus inactivation. Gently mix virus stock.
2. Make 1:250 and 1:1000 dilutions of the AAV stock(s) in DMEM + 2% FBS.  
NOTE: Each test well will receive 50 µl of the appropriate dilution. Lower or higher dilutions may be required depending on the serotype and AAV production conditions.
3. Add 50 µl of the appropriate AAV dilution to wells containing cells.
4. Incubate the assay wells at 37°C in 5% CO<sub>2</sub> for 48 hours post-transduction.
5. NOTE: To obtain an accurate titer, it is desirable to have less than 20% GFP positive cells at 48 hours post-transduction. This minimizes counting cells with multiple integration events, which would result in an underestimation of titer.


#### C. Cell harvest and analysis

1. Gently wash cells with 200 µl 1X PBS. Following the removal of PBS, immediately add 100 µl of trypsin to each well.
2. Incubate the plate at 37°C and closely monitor cell rounding and detachment.
3. After cells have detached, add 400 µl of complete growth media (e.g. DMEM + 10% FBS) to each well to inactivate the trypsin and resuspend the cells.  
NOTE: The cells should be at  $\sim 1 \times 10^6$  cells/ml. The cells can be further diluted in growth media if desired.
4. Transfer 250 µl of cell suspension from each well to separate wells in a non-treated 96-well plate (or similar culture vessel) that is compatible with your flow cytometer.  
NOTE: The optimal volume added for dilution may vary depending on the flow cytometer.



Cells can also be plated 18-24 hours before transduction, but at a lower density ( $7.5 \times 10^4$  cells/ml) to ensure 40-50% confluency at the time of transduction. If plated the day before, the cells should be trypsinized and counted again at the time of transduction because cells will likely have divided and increased in number.

Representative image of ~50% confluent HT-1080 cells:



5. Analyze for GFP expression by flow cytometry.
6. Calculate the functional titer of the AAV stock using the following equation:

$$\text{Titer (HT-1080 Transducing units/ml)} = \left[ \frac{\text{Number of target cells (Count at time of transduction)} \times [\% \text{ GFP positive cells}/100]}{\text{(Volume of AAV Stock in ml)}} \right]$$

NOTE: To determine the functional titer produced per milliliter of total culture, multiply the AAV stock titers determined above by the dilution factor at harvest (e.g. for AAV produced in suspension HEK 293 cultures and harvested as described in Section I.D, multiply AAV stock titers determined above by 1.2).

## TROUBLESHOOTING GUIDE

POOR DNA TRANSFECTION EFFICIENCY	
Problem	Solution
Incorrect vector sequence	If you do not observe expression of your target insert, verify the sequence of the transfer vector plasmid DNA.
Suboptimal <i>TransIT</i> ® Reagent:DNA ratio	Determine the best <i>TransIT</i> -VirusGEN® SELECT Reagent:DNA ratio for each cell type. Titrate the <i>TransIT</i> -VirusGEN® SELECT Reagent volume from 1-3 µl per 1 µg DNA. Refer to “Before You Start” on Page 2 for recommended starting conditions.
Suboptimal DNA concentration	Determine the DNA concentration accurately. Use plasmid DNA preps that have an A <sub>260/280</sub> absorbance ratio of 1.8-2.0. The optimal DNA concentration generally ranges between 0.5-2 µg per 1 ml of culture. Start with 2 µg DNA per 1 ml of culture. Consider testing different amounts of DNA while scaling the amount of <i>TransIT</i> -VirusGEN® SELECT accordingly.
Low-quality plasmid DNA	Use highly purified, sterile, endotoxin- and contaminant-free DNA for transfection. We recommend using Mirus MiraCLEAN® Endotoxin Removal Kit (MIR 5900) for removal of endotoxin from your DNA preparation. Alternatively, use cesium chloride gradient or anion exchange purified DNA which contains levels of endotoxin that do not harm most cells.
Cells not actively dividing at the time of transfection	Divide the culture at least 18-24 hours before transfection to ensure that the cells are actively dividing and reach optimal cell density at time of transfection. DO NOT proceed with transfection if cells are not doubling normally or are < 95% viable by trypan blue exclusion.
Transfection incubation time not optimal	Determine the optimal transfection incubation time for each cell type and experiment. Test a range of incubation times (e.g. 48-72 hours). The best post-transfection incubation time for AAV is typically 48-72 hours.
<i>TransIT</i> -VirusGEN® SELECT was not mixed properly	Warm <i>TransIT</i> -VirusGEN® SELECT Reagent to room temperature and vortex gently before each use. If <i>TransIT</i> -VirusGEN® SELECT Reagent is pre-diluted in complex formation solution, DNA should be added within 5 min. Incubating the <i>TransIT</i> -VirusGEN® SELECT Reagent in complex formation solution alone for an extended time results in reduced production of functional virus.
Disruption of transfection complex formation	After initial mixing of DNA and <i>TransIT</i> -VirusGEN® SELECT Reagent, do not agitate Reagent:DNA complexes again, e.g. do not vortex or invert immediately before adding to cultures.
Precipitate formation during transfection complex formation	During complex formation, scale all reagents according to the scaling tables provided in each section of the protocol, including: serum-free media, <i>TransIT</i> -VirusGEN® SELECT Reagent and plasmid DNA. Precipitation may be observed when excess DNA is used during complex formation. This may negatively impact transfection efficiency. To avoid precipitation when using high concentrations of DNA, increase the volume of serum-free medium during complex formation by two-fold.
Proper experimental controls were not included	To assess delivery efficiency of plasmid DNA, use Mirus <i>Label IT</i> ® Tracker™ Intracellular Nucleic Acid Localization Kit to label the target plasmid or use Mirus pre-labeled <i>Label IT</i> ® Plasmid Delivery Controls (please refer to Related Products on Page 10). To verify efficient transfection, use <i>TransIT</i> -VirusGEN® SELECT Reagent to deliver a positive control such as a luciferase, beta-galactosidase or green fluorescent protein (GFP) encoding plasmid.

## TROUBLESHOOTING GUIDE continued

HIGH CELLULAR TOXICITY	
Problem	Solution
Cell density not optimal at time of transfection	High toxicity and cell death may be observed if cells are not dense at the time of transfection. For high virus titers using <i>TransIT-VirusGEN®</i> SELECT Reagent, ensure that cell cultures are approximately $2 \times 10^6$ cells/ml (for suspension cell transfections) at the time of transfection.
Cell morphology has changed	When generating AAV with the VirusGEN® SELECT AAV Complex Formation Solution and Enhancer, cell growth may decrease. This is normal and does not adversely affect virus titers.
	Mycoplasma contamination can alter cell morphology and affect transfection efficiency. Check your cells for mycoplasma contamination. Use a fresh frozen stock of cells or use appropriate antibiotics to eliminate mycoplasma.
	A high or low cell passage number can make cells more sensitive and refractory to transfection. Maintain adherent or suspension HEK 293 cells below passage 30 for optimal recombinant virus production.
Transfection complexes not evenly distributed after complex addition to cells	Add transfection complexes while swirling the flask. If this is not possible, gently mix the culture vessel to ensure even distribution of the transfection complexes.

### RELATED PRODUCTS

- VirusGEN® GMP AAV Transfection Kit
- VirusGEN® AAV Transfection Kit
- VirusGEN® GMP LV Transfection Kit
- VirusGEN® SELECT LV Transfection Kit
- VirusGEN® LV Transfection Kit
- *TransIT*-VirusGEN® GMP Transfection Reagent
- *TransIT*-VirusGEN® SELECT Transfection Reagent
- *TransIT*-VirusGEN® Transfection Reagent
- *TransduceIT*™ Reagent
- *Label IT*® Plasmid Delivery Controls
- *Label IT*® Tracker™ Intracellular Nucleic Acid Localization Kits
- MiraCLEAN® Endotoxin Removal Kits
- Ingenio® Electroporation Solution and Kits

For details on the above-mentioned products, visit [www.mirusbio.com](http://www.mirusbio.com)



#### Reagent Agent®

Reagent Agent® is an online tool designed to help determine the best solution for nucleic acid delivery based on in-house data, customer feedback and citations.

Learn more at:  
[www.mirusbio.com/ra](http://www.mirusbio.com/ra)

Contact Mirus Bio for additional information.



Mirus Bio LLC  
5602 Research Park Blvd, Ste 210  
Madison, WI 53719  
Toll-free: 888.530.0801  
Direct: 608.441.2852  
Fax: 608.441.2849

©1996-2022 All rights reserved. Mirus Bio LLC. All trademarks are the property of their respective owners.  
For terms and conditions, visit [www.mirusbio.com](http://www.mirusbio.com)