

CHOgro® Expression Medium

Quick Reference Protocol

Instructions for MIR 6200, 6202

SDS and Certificate of Analysis available at mirusbio.com/6200



SPECIFICATIONS

Storage	Store CHOgro® Expression Medium at 4°C, protected from light.
Product Guarantee	As indicated on product label, when properly stored and handled.

CHOgro® Expression Medium is a chemically defined, hydrolysate-free and animal-origin-free medium. CHOgro® Expression Medium is formulated to support high efficiency transfection and provide high density cell growth. In addition, many suspension CHO cells (e.g. Freestyle™ CHO-S and ExpiCHO-S™) can easily grow in CHOgro® Expression Medium with minimal adaptation. CHOgro® Expression Medium is supplied in 1000 ml bottle (MIR 6200) or 10 L polybag (MIR 6202) formats and is also a component within the CHOgro® Expression System (MIR 6260) and CHOgro® High Yield Expression System (MIR 6270).

Media Preparation

Prior to use, CHOgro® Expression Medium requires the following supplementation:

Media Supplements	Per 1000 ml
L-Glutamine (200 mM stock solution, MIR 6240)	20 ml
Poloxamer 188 Solution (10% stock solution, MIR 6230)	30 ml

NOTE: Store supplemented media at 4°C, protected from light.

Adaptation of Suspension CHO cells to CHOgro® Expression Medium

From Cryopreserved Cell Stock

When bringing suspension CHO cells out of cryopreservation, use supplemented CHOgro® Expression Medium to dilute cells immediately post-thaw, typically to a density of 0.5 - 1 × 10⁵ cells/ml. Incubate cells in a shake flask at an appropriate rpm (e.g. 125 rpm for a 1.9-cm orbital throw) at 37°C in 8% CO₂. Monitor cell growth and viability daily. When viability reaches ≥ 98% and the cells are doubling every ≤ 24 hours, the cells are fully adapted.

From Ongoing Culture

If cells are cultured in an alternate media formulation, cells must be adapted to CHOgro® Expression Medium prior to transfection. For adaptation, seed cells at a density of 3 - 5 × 10⁵ cells/ml in a mix of 75% current media and 25% complete CHOgro® Medium for 2-4 passages or until the cells are doubling normally and viability is > 95%. Increase the ratio of complete CHOgro® Media in 25% increments and monitor cell health and viability as described above. Cells are fully adapted when viability reaches ≥ 98% and cells are doubling every ≤ 24 hours in 100% CHOgro® Medium. Continue to culture cells in a shake flask at an appropriate rpm (e.g. 125 rpm for 1.9-cm orbital throw) at 37°C, 8% CO₂ and monitor cell growth and viability frequently.

Maintenance of Suspension CHO cells in CHOgro® Expression Medium

For best results, subculture CHO suspension cells to a density of 1 - 3 × 10⁶ cells/ml. DO NOT allow cells to grow to a density higher than 1 × 10⁷ cells/ml or passage to a density lower than 2.5 × 10⁵ cells/ml during continuous culture. Subculture every 1-4 days to maintain the desired cell density. Monitor cell density and viability frequently (i.e. daily if possible).

For Research Use Only

CHOgro® Expression Medium is sterile-filtered and animal-origin-free.



Reagent Agent®

Reagent Agent® is an online tool designed to help determine the best solution for nucleic acid delivery based on in-house data, customer feedback and citations.

Learn more at: mirusbio.com/ra



SDS and Certificate of Analysis available at mirusbio.com/6200

©1996-2023 All rights reserved. Mirus Bio LLC. All trademarks are the property of their respective owners.
For terms and conditions, visit www.mirusbio.com

Rev1 022222