

TransIT®-Insect Transfection Reagent

Quick Reference Protocol

Instructions for MIR 6100, 6104, 6105, 6106, 6110

Full protocol, SDS and Certificate of Analysis available at mirusbio.com/6100



SPECIFICATIONS

Storage	Store TransIT®-Insect Reagent tightly capped at -20°C. <i>Before each use</i> , warm to room temperature and vortex gently.
Product Guarantee	1 year from the date of purchase, when properly stored and handled.

► PLASMID DNA TRANSFECTION PROTOCOL



Full protocol and additional documentation available at
mirusbio.com/6100

Fill in volumes below based on culture vessel used for transfection (Table 1).

A. Plate cells

1. Plate cells in ____ ml complete growth medium (per well).
For adherent cells: Plate cells at a density of $1.6\text{-}3.2 \times 10^5$ cells/ml.

For suspension cells: Plate cells at a density of $3.2\text{-}4.8 \times 10^5$ cells/ml.

2. Culture overnight. Most cell types should be approximately 80% confluent at the time of transfection.

B. Prepare TransIT®-Insect Reagent:DNA complexes

1. Warm TransIT®-Insect to room temperature and vortex gently.
2. Place ____ μ l of Grace's Insect Basal Medium in a sterile tube.
3. Add ____ μ l plasmid DNA. Mix gently by pipetting.
4. Add ____ μ l of TransIT®-Insect Reagent. Mix gently by pipetting.
5. Incubate at room temperature for 15-30 minutes.

C. Distribute complexes to cells

1. Add TransIT®-Insect:DNA complex mixture drop-wise to different areas of the well.
2. Gently rock plate for even distribution of complexes.
3. Incubate 24-72 hours.
4. Harvest cells and assay as required.

Table 1. Recommended starting conditions

Culture vessel	24-well plate	12-well plate	6-well plate
Surface area	1.9 cm ²	3.8 cm ²	9.6 cm ²
Complete growth medium	0.5 ml	1 ml	2.5 ml
Serum-free medium	50 μ l	100 μ l	250 μ l
DNA (1 μ g/ μ l stock)	0.5 μ l	1 μ l	2.5 μ l
TransIT®-Insect Reagent	1 μ l	2 μ l	5 μ l

► Transfection Optimization

Determine the best TransIT®-Insect Reagent:DNA ratio for each cell type. Start with 2 μ l of TransIT®-Insect Reagent per 1 μ g of DNA. Vary the concentration of TransIT®-Insect Reagent from 1-4 μ l per 1 μ g DNA to find the optimal ratio.

For additional optimization tips, see [full protocol](#). Cell-type-specific recommendations available on [Reagent Agent](#) (mirusbio.com/ra).

► NOTES



Reagent Agent[®]

Reagent Agent[®] is an online tool designed to help determine the best solution for nucleic acid delivery based on in-house data, customer feedback and citations.

Learn more at: mirusbio.com/ra

©1996-2024 All rights reserved. Mirus Bio LLC. All trademarks are the property of their respective owners.

For terms and conditions, visit www.mirusbio.com

Rev0 01282022

Mirus Bio LLC

www.mirusbio.com | techsupport@mirusbio.com | U.S. Toll Free: 844.MIRUSBIO (844.647.8724) | Direct: +1.608.441.2852